Wellness Microbreak

A microbreak is a break you take every 30 to 60 minutes. Stop your current work activity, drink some water, change position, do a stretch and change posture. It can only take a minute. Then, reposition yourself and resume work.

## Hydration is important for wellness.

Drink approximately eight, 8-ounce glasses of water per day. Tea, coffee and cola drinks pull water out of you; they are diuretics so they don't count.

These exercises are to be done as gentle stretches; they are intended for healthy adults and for educational purposes, not as medical advice. If you have any medical conditions or have had injuries to your neck, shoulders, spine, or arms, check with your health care professional before doing these exercises. If any of the exercises cause pain, STOP. Do not continue stretching through the pain.

## **Gentle Wrist & Hand Movements**

CAUTION: Do not try these stretches if you have numbness in your hands or have been diagnosed with carpal tunnel syndrome or tendonitis at the wrist.





### Wrist Stretch

- Hands together with palms facing
- Keep palms flat together
- Raise elbows, drop wrists, roll shoulders back



## **Carpal Tunnel Stretch**

- Arm against side, elbow bent, hand held out in front of you, palm up
- Using fingers of other hand, apply slow, gentle downward pressur on outstretched palm, bending hand back slowly
- STOP when you feel a gentle pull
- Optional: Repeat exercise with elbow straight to stretch biceps

# **Hand & Wrist Series**



Position 1 Wrist in neutral, fingers & thumb in flexion



Wrist in neutral, fingers & thumb extended


**Eye Relaxation** 



- Focus on an object at least 20 feet away
- Repeat 3 times




- Shape hands into shallow cups
- Place hands lightly over eyes • Hold them there for one minute



• Look up, right, down & left 3 times



### **Rotate Eyes**



### **Gentle Neck Movements**




### **Head Tilt**

- Sit upright with your spine relaxed (neutral)
- Lower left shoulder
- Tilt head to right
- Hold 10 seconds · Repeat on opposit



### **Head Rotation**

- Sit upright with your spine relaxed (neutral)
- Turn head slowly to right
- Repeat on opposite side



Position 4

spread apart

Same as Position 3

with fingers & thumb

## **Neck Retraction**

- Face forward
- Keep head & ears level
- Slide head forward & back
- Repeat 5-10 times



Position 6

Same as Position 5 with other hand gently stretching thumb

Position 5 Wrist in neutral, fingers & thumb same as Position 4



**Shoulder Rolls** 

**Gentle Lower & Mid Back Movements** 

Reverse Slouch

• Sit relaxed, hands on knees

Mid Back Realignment

**Abdominal Strengthening** 

**Gentle Shoulder & Upper Back Movements** 

**Upper Back** 

Hands behind upper back

High Reach Stretch

Spread fingers

Bend forward, elbows toward floor

• Unwind, lower back first, head last

• Raise arms reaching above head

• Hold for 5 seconds, relax & repeat 5 times

• Reach as high as you can

• Curl head forward, drop arms

• Slowly let hands & body curl toward floor

Slowly uncurl, from lower back up to head

• Raise one arm beside & touching ear

• Bend this arm at elbow so hand is behind head

• Opposite arm is down & bent behind back; hand

reaching toward mid back (trying to touch other hand)

• Tighten abdominal muscles while seated or standing


• Hold for count of 10, release & repeat 5-10 times

• If possible, support lower back with hands & arch back gently

- Sit comfortably; back upright & relaxed, feet flat on the floor
- Gently roll shoulders backward; relax & repeat 5 times
- Reverse directions and repeat above steps, if needed

# Shoulder Squeeze

- Sit upright with your spine relaxed (neutral)
- Elbows at shoulder height
- Move arms back; squeeze shoulder blades together & hold
- Slide head back over shoulders



### **Shoulder Shrug**

- Sit upright with your spine relaxed (neutral)
- Shrug shoulders up & hold • Relax & repeat 5 times



Position 3

Wrist, fingers &

thumb extended back

Kathy Davis, PhD; Paul Dougherty, DC; Marjorie Eskay-Auerbach, MD; Helen K. Maher, PhD, RN, COHN-S; Catherine M. Pepler RN, BS, MBA, COHN-S/CM, FAAOHN and Kimberly Templeton, MD. Special thanks to Mary Gene Ryan, MPH, BSN, RN, COHN-S/SM, FAAOHN and MGRyan & Co., Inc. for their collaboration and generosity.





